International Prostate Symptom Score (I-PSS)

PATIENT NAME

Please put a rating next to each of the questions shown below. Then add up the numbers and write the total at the bottom. For the following questions the rating is determined as below:

- 0 = Not at all
- 3 = About half the time
- 1 = Less than 1 time in 52 = Less than half the time
- 4 = More than half the time 5 = Almost always
- 1. Incomplete Emptying

Over the past month, how often have you had a sensation of not emptying your bladder completely after you finish urinating?

..... 2. Frequency

Over the past month, how often have you had to urinate again less than two hours after you have finished urinating?

...... 3. Intermittency

Over the past month, how often have you found you stopped and started several times when you urinated?

..... 4. Urgency

Over the past month, how often have you found it difficult to postpone urination?

..... 5. Weak Stream

Over the last month, how often have you had a weak urinary stream?

..... 6. Straining

Over the past month, how often have you had to push or strain to begin urination?

...... 7. Nocturia

Over the past month how many times did you most typically get up each night to urinate from the time you went to bed until the time you got up in the morning?

- o = None 1 = Once
- 2 = Twice
- Total Score
- 3 = Three times 4 = Four times
- 5 = Five times or more

...... 8. Quality of Life due to Urinary Symptoms:

If you were to spend the rest of your life with your symptoms just the way they have been during the last week, how would you feel about that?

- o = Delighted
- = Pleased

1

2

3

4

6

- Mostly satisfied
- = Mixed (about equally satisfied and dissatisfied)
- Mostly dissatisfied
- 5 = Unhappy
 - = Terrible

The I-PSS is based on the answers to seven questions concerning urinary symptoms. Each question is assigned points from 0 to 5 indicating increasing severity of the particular symptom. The total score can therefore range from 0 to 35 (asymptomatic to very symptomatic).

Although there are presently no standard recommendations into grading patients with mild, moderate or severe symptoms, patients can be tentatively classified as follows:

0-7 = mildly symptomatic;
8-19 = moderately symptomatic;
20-35 = severely symptomatic.

The International Consensus Committee (ICC) recommends the use of only a single question to assess the patient's quality of life. The answers to this question range from "delighted" to "terrible" or o to 6. Although this single question may or may not capture the global impact of BPH symptoms on quality of life, it may serve as a valuable starting point for doctor-patient conversation.